

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
7 July 2005 (07.07.2005)

PCT

(10) International Publication Number
WO 2005/060768 A1

(51) International Patent Classification⁷: **A23L 1/305**,
1/29, A61K 31/401

FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO,
SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN,
GQ, GW, ML, MR, NE, SN, TD, TG).

(21) International Application Number:
PCT/EP2004/013787

Declarations under Rule 4.17:

(22) International Filing Date: 3 December 2004 (03.12.2004)

— as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW, ARIPO patent (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Date:
03029505.9 20 December 2003 (20.12.2003) EP

(71) Applicant (for all designated States except US): NESTEC S.A. [CH/CH]; Avenue Nestlé 55, CH-1800 Vevey (CH).

(72) Inventors; and

(75) Inventors/Applicants (for US only): SMOLA, Hans [DE/DE]; Ubiering 14-16, 50678 Koeln (DE). NEPO-MUCENO, Gilberto [BR/CH]; Chisenmattweg 8, CH-3510 Konolfingen (CH).

(74) Agent: DIXON, Sarah; Avenue Nestlé 55, CH-1800 Vevey (CH).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,

— as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW, ARIPO patent (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: NUTRITIONAL COMPOSITION FOR WOUND HEALING

(57) Abstract: A nutritional composition for promoting wound healing comprises a protein source, a lipid source and a carbohydrate source wherein no more than 1.8 % of the total calories of the composition derive from arginine and wherein the protein source includes proline in an amount of at least 3 % of the total calories of the composition. The composition may be administered orally and is particularly suitable for the amelioration of pressure ulcers although it may also be used with advantage in the nutritional management of acute wounds including pre and post surgery.

WO 2005/060768 A1

Nutritional Composition for Wound Healing

Field of the invention

- 5 This invention relates to a nutritional composition for promoting wound healing, particularly the healing of chronic wounds such as pressure ulcers (decubitus).

Background of the invention

- 10 In normal wound healing, there are three phases which overlap to some extent. Briefly, the first phase is inflammation in which the clot forms and stops the bleeding from blood vessels followed by extravasation of mononuclear blood cells which clean the wound and remove debris. The next phase is the granulation phase in which fibroblasts proliferate and accumulate in the wound and produce collagen to assist in
15 wound closure. This phase is characterised by high metabolic activity. Finally, epithelial cells begin to cover the wound.

- Delayed or impaired wound healing is a problem for health care professionals and patients as it results in increased treatment times and stays in healthcare facilities and distress to patients. The process of wound healing can be interrupted in any of the
20 phases described above as a result of factors such as infection or malnutrition. The pressure ulcers which frequently afflict elderly and bed-ridden patients are a particular concern and these categories of patients are often found to be suffering from malnutrition. Indeed, all patients with acute or chronic wounds exhibit
25 increased nutritional requirements, displaying a need for increased nutrients and energy as compared with individuals not challenged by such metabolic stresses. If these patients are malnourished before suffering wounds, the wounds may simply fail to heal.

- 30 In recent years, much attention has focused on the role of arginine in wound healing. This is discussed for example in USP 5,053,387 which discloses an enteral nutritional formulation in which 1 to 3% of the total energy intake is preferably provided by arginine. Similarly, EP 960 572 A discloses a nutritional composition suitable for the treatment and prevention of pressure ulcers which includes arginine as well as large
35 amounts of vitamins C and E. The role of arginine is also discussed in USP 5,733,884 which discloses a method of providing nutrition to a patient with an acute

or chronic wound using a composition in which at least 2% of the energy is provided by arginine and the same amount by proline. This patent hypothesises that arginine and proline have a synergistic effect in enhancing wound healing. Commercially, there are a number of products marketed as suitable for promoting wound healing on the basis that they contain high levels of arginine including CUBITAN® and ARGINAID®.

An adequate supply of arginine is clearly relevant to the wound healing process. However, arginine is also a precursor for the formation of nitric oxide which acts as a vasodilator and enhances growth hormone secretion. It is not desirable for critically ill individuals to be exposed to high amounts of nitric oxide and yet this will inevitably happen if such individuals receive nutritional supplements containing high levels of arginine – see, for example L. Cynober, Curr Opin Clin Nutr Metab Care. 6:189-93 2003. Moreover, it is quite likely that a high proportion of elderly, bedridden or critically ill patients at risk of developing pressure sores will also suffer from conditions for which high levels of nitric oxide are contra-indicated (J. Takala et al., N Engl J Med 341:785-792 1999).

Summary of the Invention

In a first aspect, the present invention provides a nutritional composition for promoting wound healing comprising a protein source, a lipid source and a carbohydrate source wherein no more than 1.8% of the total calories of the composition derive from arginine and wherein the protein source includes proline in an amount of at least 3% of the total calories of the composition.

In a second aspect, the present invention provides a method of providing nutritional support to a patient with an acute or chronic wound comprising the step of administering a therapeutically effective amount of a nutritional composition comprising a protein source, a lipid source and a carbohydrate source wherein no more than 1.8% of the total calories of the composition derive from arginine and wherein the protein source includes proline in an amount of at least 3% of the total calories of the composition.

In a third aspect, the present invention provides the use of a protein source, a lipid source and a carbohydrate source for the manufacture of a therapeutic formulation for promoting wound healing wherein no more than 1.8% of the total calories of the formulation derive from arginine and wherein the protein source includes proline in an amount of at least 3% of the total calories of the formulation.

Detailed Description of the Invention

Although the inflammatory phase of the wound healing process described above is critical, the present inventor believes that from a therapeutic/nutritional approach, attempts to modulate this phase carry high risks and that the granulation phase offers better potential for nutritional intervention. In this phase, new connective tissue is synthesised and more than 80% of this tissue is composed of collagen. Collagen is rich in the amino acids proline (about 22%) and glycine (about 33%) and the presence of these amino acids is rate limiting for collagen formation, that is to say, collagen cannot be efficiently formed if they are not available in sufficient quantity. However, the normal diet contains only about 3% in total of these amino acids and it will be appreciated that individuals who have suffered wounds may ingest even less of them, particularly as proline is not generally regarded as an essential dietary amino acid. In the case of individuals suffering from malnutrition for whatever reason, these shortages may be particularly pronounced. The composition of the present invention is therefore supplemented with proline in a quantity sufficient to facilitate collagen synthesis. It is particularly suitable for the amelioration of pressure ulcers but may also be used in the management of acute wounds including before and after surgery.

The composition of the present invention does not need to be supplemented with arginine - of course some arginine is likely to be present as part of the protein source. However, it is widely believed that arginine also has a role in the inflammatory phase of wound healing and, for this reason, the composition of the present invention is preferably supplemented with small amounts of arginine subject always to the requirement that arginine must account for no more than 1.8% of the total calories of the composition..

The composition of the present invention contains sources of protein, lipids and carbohydrate and may be administered orally or enterally. The composition preferably provides about 1.25 kcal/ml.

- 5 Protein is essential to healing as tissue damage results in a catabolic response that includes a requirement for a larger proportion of total calories as protein than is required by the general population. Research suggests that enteral fortification employing large quantities of protein can accelerate the synthesis of visceral proteins and so the protein source of the present invention preferably constitutes at least 25%
10 of the total energy content of the composition, more preferably at least 28%.

- A variety of different protein sources may be used including intact protein sources such as casein or whey as well as hydrolysed proteins, free amino acids and even mixtures of intact and hydrolysed proteins and/or free amino acids, in each case
15 supplemented with free proline and, optionally, free arginine. Preferably, the protein source of the present invention is selected to yield the highest amount of proline in the proteins so as to minimise the amount that needs to be added as the free amino acid.

- 20 Preferably, proline constitutes at least 3.5% of the calories of the composition of the present invention. At this level of contribution to total calories, the composition will need to be supplemented by about 3.0% (by weight of the protein source) proline.

- The total calories/gram of nitrogen of the composition of the invention is preferably
25 about 160:1. The total non-protein calories/ gram of nitrogen is preferably about 110:1.

- The composition of the present invention also includes a lipid source. Lipids or fats are the primary source of stored energy in the body and energy from fat metabolism is
30 used in all normal cell functions. As far as wound healing is concerned, fat metabolism results in the formation of prostaglandins and other regulators of the inflammatory process. The lipid source used in the present invention preferably constitutes about 20% of the total energy content of the composition. Of this 20%, preferably about 8% is constituted by mono- and di-glycerides of fatty acids. The

ratio of n-6 to n-3 fatty acids is preferably between 4:1 and 10:1, more preferably about 7:1.

The composition of the present invention also includes a carbohydrate source. Glucose is the primary fuel for cellular metabolism of many tissues including leucocytes, macrophages and fibroblasts all of which are involved in the wound healing process. Glucose is needed to meet the specific metabolic demands of wound healing. The carbohydrate source used in the present invention preferably constitutes about 50% of the total energy content of the composition. Suitable sources of carbohydrate are maltodextrin and sucrose. Preferably, the carbohydrate source is substantially free of lactose.

Vitamins, minerals and trace elements are also important in the wound healing process. Preferably, the composition of the present invention at least complies with the compositional criteria set out in Directive 1999/21/EC on Dietary Foods for Special Medical Purposes as regards these micronutrients. However, certain micronutrients are particularly important for wound healing and therefore the composition of the present invention preferably contains more than the recommended minimum levels of vitamins C and E, manganese, zinc and selenium.

A liquid, ready to use composition according to the present invention will now be given by way of example:-

Example 1

Caloric density	1.25g/ml	
Protein	30% of kcal	
of which (by weight):-	sodium caseinate	50%
	milk protein concentrate	45%
	free L-proline	3%
	free L-arginine	2%
total L-proline	12.4% of protein source	
total L-arginine	5.0% of protein source	
Caloric contribution of total proline		3.7%
Caloric contribution of total arginine		1.5%

	Lipids	20% of kcal	
	of which	rapeseed oil	35%
		corn oil	34%
5		soya oil	20%
		mono and di-glycerides of fatty acids	8%
		milk fat	3%
	n-6:n-3	7.2:1	
10	Carbohydrate	50% of kcal	
	of which	corn syrup	52%
		sucrose	43%
		starch	3%
		lactose	2%
15	Vitamin C	125mg/100ml	
	Vitamin E	7.5mg α -tocopherol equivalents/100ml	
	Manganese	1.9mg/100ml	
	Zinc	3.7mg/100ml	
	Selenium	19 μ g/100ml	
20	Osmolarity	470 mosm/Kg water	
	Water	80.3%	
	Density	1.087g/ml	
	Total cal/g nitrogen	160:1	
25	Non-protein cal/g nitrogen	110:1	

As will be appreciated from the foregoing description, the composition will also contain other micronutrients of the type conventionally found in enteral compositions in accordance with EC Directive 1999/21/EC as well as flavourings such as coffee or vanilla, emulsifiers, thickeners and stabilizers of the type conventionally found in enteral compositions.

The nutritional composition may be produced by conventional methods. For example, the protein source and the lipid source are dissolved in water, preferably water which has been subjected to reverse osmosis, to form a liquid mixture.

Emulsifiers may be dissolved in the lipid source prior to blending if desired. Preferably, a food grade emulsifier from a vegetable source is used.

5 The temperature of the water is conveniently about 50°C to about 80°C to aid dispersal of the ingredients. Commercially available liquefiers may be used to form the liquid mixture. Preferably, pH of the liquid mixture is adjusted to about 6.3 to 7 with food grade hydroxides.

10 After preparation of the liquid mixture, the carbohydrate source is added together with other easily dissolvable ingredients including, for example, vitamins, minerals, flavourings and colorants.

15 The liquid mixture may then be thermally treated to reduce bacterial loads (pasteurized). This may be carried out by steam injection or by heat exchanger; for example a plate heat exchanger.

20 If a shelf-stable liquid composition is required, an ultra heat treatment (UHT) is preferably conducted after pre-heating to 50-85°C. For example, an indirect UHT treatment may be conducted at 140-155°C for 5-8s, in a tube heat exchanger. The liquid mixture may then be cooled to about 60°C to about 85°C; for example by flash cooling. The liquid mixture is then homogenized and the resulting homogenised milky liquid may be aseptically filled into suitable containers such as 200 ml cups for oral feeding. Aseptic filling of the containers may be carried out by cooling the liquid mixture.

25 If a powdered, reconstitutable formula is required, the homogenised mixture can be evaporated and dried to powder; for example by spray drying. Conventional procedures may be used.

30

Experimental Example

35 Normal human fibroblasts were trypsinised and seeded in 12 well plates at a density of 10,000 cells/cm². When confluent, the cells were transferred to a culture medium with an amino acid distribution and concentrations designed to mimic those in human

serum as closely as possible. The cell cultures were divided into two categories, a control culture in which the culture medium contained 0.201 mM proline and an experimental sample in which the culture medium contained 0.592 mM proline. After 24 hours fibroblast-conditioned medium containing 100 microgram/ml beta-aminopropionitrile to prevent cross-linking of collagen molecules in the cultures was collected. The conditioned medium was dotblotted to a nitrocellulose membrane and probed for collagen type I content with a polyclonal immune-absorbed antibody. The value shown for the proline-supplemented samples is relative to the controls set at 100%.

Sample	% of control value
Control (0.201 mM Proline)	100%
Proline-supplemented (0.502 mM Proline)	150% \pm 21.9%

This experimental example shows that human fibroblasts respond to proline supplementation with a 50% increase in collagen synthesis. In this proline-supplemented medium, increased collagen synthesis is independent of the addition of growth factors or other mediators stimulating collagen transcription. It indicates an increased substrate requirement for efficient collagen synthesis.

Claims

1. A nutritional composition for promoting wound healing comprising a protein source, a lipid source and a carbohydrate source wherein no more than 1.8% of the total calories of the composition derive from arginine and wherein the protein source includes proline in an amount of at least 3% of the total calories of the composition..
2. A nutritional composition according to Claim 1 wherein at least 3.5% of the total calories of the composition derive from proline.
3. A nutritional composition according to claim 1 or 2 wherein 1.5% of the total calories of the composition derive from arginine.
4. A nutritional composition according to any preceding claim wherein the protein source constitutes at least 28% of the total calories of the composition.
5. A nutritional composition according to any preceding claim which composition has an energy density of about 1.25kcal/ml.
6. A method of providing nutritional support to a patient with an acute or chronic wound comprising the step of administering a therapeutically effective amount of a nutritional composition comprising a protein source, a lipid source and a carbohydrate source wherein no more than 1.8% of the total calories of the composition derive from arginine and wherein the protein source includes proline in an amount of at least 3% of the total calories of the composition.
7. A method according to claim 6 wherein at least 3.5% of the total calories of the composition derive from proline.
8. A method according to claim 6 or 7 wherein 1.5% of the total calories of the composition derive from arginine.

9. The use of a protein source a lipid source and a carbohydrate source for the manufacture of a therapeutic formulation for promoting wound healing, wherein no more than 1.8% of the total calories of the composition derive from arginine and wherein the protein source includes proline in an amount of at least 3% of the total calories of the formulation.
- 5
10. The use according to claim 9 wherein at least 3.5% of the total calories of the formulation derive from proline.
- 10
11. The use according to claim 9 or 10 wherein 1.5% of the total calories of the formulation derive from arginine

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP2004/013787A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 A23L1/305 A23L1/29 A61K31/401

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 7 A23L A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

EP0-Internal, WPI Data, PAJ, FSTA, BIOSIS

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5 384 308 A (HENKIN R I) 24 January 1995 (1995-01-24) claims 1,2 column 2, line 15 - column 3, line 68 column 5, lines 16-43 column 8, lines 9-21 -----	1-11
X	DE 102 21 403 A (KYBERG PHARMA VERTRIEBS GMBH &) 4 December 2003 (2003-12-04) claims 1,2,12,13,22 -----	1-11
Y	EP 0 764 405 A (CLINTEC NUTRITION CO) 26 March 1997 (1997-03-26) example 2 page 2, lines 26-31,53,54 page 3, lines 36-41 ----- -/-	1-11

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
 "E" earlier document but published on or after the International filing date
 "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
 "O" document referring to an oral disclosure, use, exhibition or other means
 "P" document published prior to the International filing date but later than the priority date claimed

"T" later document published after the International filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
 "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the International search 8 March 2005	Date of mailing of the International search report 18/03/2005
Name and mailing address of the ISA European Patent Office, P.B. 5618 Patentlaan 2 NL - 2280 HV Rijswijk Tel: (+31-70) 340-3340, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016	Authorized officer Tallgren, A

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP2004/013787

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 01/78532 A (NESTLE SA ;BOZA JULIO (CH); BALLEVRE OLIVIER (CH); FINOT PAUL ANDR) 25 October 2001 (2001-10-25) claims 2,4,9,16; examples 1,2 page 4, lines 1-9,30-32 page 5, line 28 - page 7, line 17 -----	1-11
Y	WO 03/075903 A (UNIVERSITEIT LEIDEN ;MATYSIK JOERG (NL); ALIA (NL); BACKENDORF CLA) 18 September 2003 (2003-09-18) claims 1,2,6-9,22,30 page 3, lines 10-20 page 8, lines 7-20 -----	1-11
Y	EP 1 161 945 A (SOLARTIUM ESTABLISHMENT) 12 December 2001 (2001-12-12) claims 1,6 page 2, lines 5-8,21-24 page 3, lines 15-31 -----	1-11
Y	WO 03/013487 A (DIOGUARDI FRANCESCO SAVERIO ;CONTI FRANCO (IT); PROFESSIONAL DIETE) 20 February 2003 (2003-02-20) claims 1,7 page 3, lines 2-12 page 4, lines 18-35 -----	1-11

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No.

PCT/EP2004/013787

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 5384308	A	24-01-1995	US 5788967 A	04-08-1998
DE 10221403	A	04-12-2003	DE 10221403 A1	04-12-2003
EP 0764405	A	26-03-1997	US 5714472 A	03-02-1998
			AT 227520 T	15-11-2002
			DE 69624768 D1	19-12-2002
			DE 69624768 T2	02-10-2003
			DK 764405 T3	10-02-2003
			EP 0764405 A2	26-03-1997
			ES 2185744 T3	01-05-2003
			PT 764405 T	31-03-2003
WO 0178532	A	25-10-2001	CA 2404005 A1	25-10-2001
			CN 1422125 A	04-06-2003
			WO 0178532 A1	25-10-2001
			EP 1274322 A1	15-01-2003
			HU 0300299 A2	28-07-2003
			JP 2003530411 T	14-10-2003
			MX PA02010058 A	10-03-2003
			NO 20024743 A	03-12-2002
			PL 358876 A1	23-08-2004
			US 2004096478 A1	20-05-2004
WO 03075903	A	18-09-2003	EP 1342474 A1	10-09-2003
			AU 2003214719 A1	22-09-2003
			WO 03075903 A2	18-09-2003
EP 1161945	A	12-12-2001	IT MI20001257 A1	07-12-2001
			EP 1161945 A2	12-12-2001
			US 2002013359 A1	31-01-2002
WO 03013487	A	20-02-2003	IT T020010804 A1	10-02-2003
			CA 2455776 A1	20-02-2003
			EP 1414431 A2	06-05-2004
			WO 03013487 A2	20-02-2003
			JP 2005501068 T	13-01-2005
			US 2004192756 A1	30-09-2004